

GTPASE ASSOCIATED PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of GTPase associated proteins and to the use of these sequences in the diagnosis, treatment, and prevention of cell proliferative, autoimmune/inflammatory, and immune system disorders.

5

BACKGROUND OF THE INVENTION

Guanine nucleotide binding proteins (GTP-binding proteins) participate in a wide range of regulatory functions in all eukaryotic cells, including metabolism, cellular growth, differentiation, signal transduction, cytoskeletal organization, and intracellular vesicle transport and secretion. In higher organisms they are involved in signaling that regulates such processes as the immune response (Aussel, C. et al (1988) J. Immunol. 140:215-220), apoptosis, differentiation, and cell proliferation including oncogenesis (Dhanasekaran, N. et al. (1998) Oncogene 17:1383-1394). Exchange of bound GDP for GTP followed by hydrolysis of GTP to GDP provides the energy that enables GTP-binding proteins to alter their conformation and interact with other cellular components. The superfamily of GTP-binding proteins consists of several families and may be grouped as translational factors, heterotrimeric GTP-binding proteins involved in transmembrane signaling processes (also called G-proteins), and low molecular weight GTP-binding proteins including the proto-oncogene Ras proteins and products of rab, rap, rho, rac, smg21, smg25, YPT, SEC4, and ARF genes, and tubulins (Kaziro, Y. et al. (1991) Ann. Rev. Biochem. 60:349-400). In all cases, the GTPase activity is regulated through interactions with other proteins.

GTP-binding proteins involved in protein biosynthesis include initiation factor 2 (IF-2), elongation factor 2 (EF-Tu), and elongation factor G (EF-G), observed in prokaryotes; and initiation factor 2 (eIF-2), elongation factor 1 α (EF-1 α) and elongation factor 2 (EF-2) observed in eukaryotes (Kaziro, supra). IF-2 promotes the GTP-dependent binding of the tRNA to the small subunit of the ribosome, the step that initiates protein translation. Similarly, elongation factors promote the binding of tRNA and GTP and the displacement of GDP after hydrolysis as protein biosynthesis proceeds.

Heterotrimeric GTP-binding proteins are composed of 3 subunits (α , β and γ) which, in their inactive conformation, associate as a trimer at the inner face of the plasma membrane. G α binds GDP or GTP and contains the GTPase activity. The $\beta\gamma$ complex enhances binding of G α to a receptor. G γ is necessary for the folding and activity of G β . (Neer, E.J. et al. (1994) Nature 371:297-300.) Multiple homologs of each subunit have been identified in mammalian tissues, and different combinations of subunits have specific functions and tissue specificities. (Spiegel, A.M. (1997) J.

Inher. Metab. Dis. 20:113-121.) G protein activity is triggered by seven-transmembrane cell surface receptors (G-protein coupled receptors) which respond to lipid analogs, amino acids and their derivatives, peptides, cytokines, and specialized stimuli such as light, taste, and odor. Activation of the receptor by its stimulus causes the replacement of the G protein-bound GDP with GTP. G α -GTP dissociates from the receptor/ $\beta\gamma$ complex and each of these separated components can interact with and regulate downstream effectors. The signaling stops when G α hydrolyzes its bound GTP to GDP and reassociates with the $\beta\gamma$ complex (Neer, supra).

The alpha subunits of heterotrimeric G proteins can be divided into four distinct classes. The α -s class is sensitive to ADP-ribosylation by pertussis toxin which uncouples the receptor:G-protein interaction. This uncoupling blocks signal transduction to receptors that decrease cAMP levels which normally regulate ion channels and activate phospholipases. The inhibitory α -I class is also susceptible to modification by pertussis toxin which prevents α -I from lowering cAMP levels. Two novel classes of α subunits refractory to pertussis toxin modification are α -q, which activates phospholipase C, and α -12, which has sequence homology with the *Drosophila* gene concertina and may contribute to the regulation of embryonic development (Simon, M.I. (1991) Science 252:802-808).

The mammalian G β and G γ subunits, each about 340 amino acids long, share more than 80% homology. The G β subunit (also called transducin) contains seven repeating units, each about 43 amino acids long. The activity of both subunits may be regulated by other proteins such as calmodulin and phosducin or the neural protein GAP 43 (D. Clapham and E. Neer, 1993, Nature 365:403-406). The β and γ subunits are tightly associated. The β subunit sequences are highly conserved between species, implying that they perform a fundamentally important role in the organization and function of G-protein linked systems (Van der Voorn L. (1992) Febs. Lett. 307 (2):131-134). They contain seven tandem repeats of the WD-repeat sequence motif, a motif found in many proteins with regulatory functions. WD-repeat proteins contain from four to eight copies of a loosely conserved repeat of approximately 40 amino acids which participates in protein-protein interactions. Mutations and variant expression of β transducin proteins are linked with various disorders. Mutations in LIS1, a subunit of the human platelet activating factor acetylhydrolase, cause Miller-Dieker lissencephaly. RACK1 binds activated protein kinase C, and RbAp48 binds retinoblastoma protein. CstF is required for polyadenylation of mammalian pre-mRNA in vitro and associates with subunits of cleavage-stimulating factor. Defects in the regulation of β -catenin contribute to the neoplastic transformation of human cells. The WD40 repeats of the human F-box protein β TrCP mediate binding to β -catenin, thus regulating the targeted degradation of β -catenin by

ubiquitin ligase (Neer, *supra*; Hart, M. et al (1999) *Curr. Biol.* 9:207-210). The γ subunit primary structures are more variable than those of the β subunits. They are often post-translationally modified by isoprenylation and carboxyl-methylation of a cysteine residue four amino acids from the C-terminus; this appears to be necessary for the interaction of the $\beta\gamma$ subunit with the membrane and with other GTP-binding proteins. The $\beta\gamma$ subunit has been shown to modulate the activity of isoforms of adenylyl cyclase, phospholipase C, and some ion channels. It is involved in receptor phosphorylation via specific kinases, and has been implicated in the p21ras-dependent activation of the MAP kinase cascade and the recognition of specific receptors by GTP-binding proteins. (Clapham and Neer, *supra*).

G-proteins interact with a variety of effectors including adenylyl cyclase (Clapham and Neer, *supra*). The signaling pathway mediated by cAMP is mitogenic in hormone-dependent endocrine tissues such as adrenal cortex, thyroid, ovary, pituitary, and testes. Cancers in these tissues have been related to a mutationally activated form of a $G\alpha$, known as the gsp (Gs protein) oncogene (Dhanasekaran, *supra*). Another effector is phosducin, a retinal phosphoprotein, which forms a specific complex with retinal $G\beta$ and $G\gamma$ ($G\beta\gamma$) and modulates the ability of $G\beta\gamma$ to interact with retinal $G\alpha$ (Clapham and Neer, *supra*).

Irregularities in the GTP-binding protein signaling cascade may result in abnormal activation of leukocytes and lymphocytes, leading to the tissue damage and destruction seen in many inflammatory and autoimmune diseases such as rheumatoid arthritis, biliary cirrhosis, hemolytic anemia, lupus erythematosus, and thyroiditis. Abnormal cell proliferation, including cyclic AMP stimulation of brain, thyroid, adrenal, and gonadal tissue proliferation is regulated by G proteins. Mutations in $G\alpha$ subunits have been found in growth-hormone-secreting pituitary somatotroph tumors, hyperfunctioning thyroid adenomas, and ovarian and adrenal neoplasms (Meij, J.T.A. (1996) *Mol. Cell. Biochem.* 157:31-38; Aussel, *supra*).

LMW GTP-binding proteins are GTPases which regulate cell growth, cell cycle control, protein secretion, and intracellular vesicle interaction. They consist of single polypeptides which, like the alpha subunit of the heterotrimeric GTP-binding proteins, are able to bind to and hydrolyze GTP, thus cycling between an inactive and an active state. LMW GTP-binding proteins respond to extracellular signals from receptors and activating proteins by transducing mitogenic signals involved in various cell functions. The binding and hydrolysis of GTP regulates the response of LMW GTP-binding proteins and acts as an energy source during this process (Bokoch, G. M. and Der, C. J. (1993) *FASEB J.* 7:750-759).

At least sixty members of the LMW GTP-binding protein superfamily have been identified

and are currently grouped into the ras, rho, arf, sar1, ran, and rab subfamilies. Activated ras genes were initially found in human cancers, and subsequent studies confirmed that ras function is critical in determining whether cells continue to grow or become differentiated. Ras1 and Ras2 proteins stimulate adenylate cyclase (Kaziro, supra), affecting a broad array of cellular processes. Stimulation of cell surface receptors activates Ras which, in turn, activates cytoplasmic kinases. These kinases translocate to the nucleus and activate key transcription factors that control gene expression and protein synthesis (Barbacid, M. (1987) Ann. Rev Biochem. 56:779-827, Treisman, R. (1994) Curr. Opin. Genet. Dev. 4:96-98). Other members of the LMW GTP-binding protein superfamily have roles in signal transduction that vary with the function of the activated genes and the locations of the GTP-binding proteins that initiate the activity. Rho GTP-binding proteins control signal transduction pathways that link growth factor receptors to actin polymerization, which is necessary for normal cellular growth and division. The rab, arf, and sar1 families of proteins control the translocation of vesicles to and from membranes for protein processing, localization, and secretion. Vesicle- and target- specific identifiers (v-SNAREs and t-SNAREs) bind to each other and dock the vesicle to the acceptor membrane. The budding process is regulated by the closely related ADP ribosylation factors (ARFs) and SAR proteins, while rab proteins allow assembly of SNARE complexes and may play a role in removal of defective complexes (J. Rothman and F. Wieland (1996) Science 272:227-234). Ran GTP-binding proteins are located in the nucleus of cells and have a key role in nuclear protein import, the control of DNA synthesis, and cell-cycle progression (Hall, A. (1990) Science 249:635-640; Barbacid, M. (1987) Ann. Rev Biochem. 56:779-827; Ktistakis, N. (1998) BioEssays 20:495-504; and Sasaki, T. and Takai, Y. (1998) Biochem. Biophys. Res. Commun. 245:641-645).

The cycling of LMW GTP-binding proteins between the GTP-bound active form and the GDP-bound inactive form is regulated by additional proteins. Guanosine nucleotide exchange factors (GEFs) increase the rate of nucleotide dissociation by several orders of magnitude, thus facilitating release of GDP and loading with GTP. The best characterized is the mammalian homologue of the Drosophila Son-of-Sevenless protein. Certain Ras-family proteins are also regulated by guanine nucleotide dissociation inhibitors (GDIs), which inhibit GDP dissociation. The intrinsic rate of GTP hydrolysis of the LMW GTP-binding proteins is typically very slow, but it can be stimulated by several orders of magnitude by GTPase-activating proteins (GAPs) (Geyer, M. and Wittinghofer, A. (1997) Curr. Opin. Struct. Biol. 7:786-792). Both GEF and GAP activity may be controlled in response to extracellular stimuli and modulated by accessory proteins such as RalBP1 and POB1. Mutant Ras-family proteins, which bind but can not hydrolyze GTP, are permanently activated, and cause cell proliferation or cancer, as do GEFs that inappropriately activate LMW GTP-binding proteins, such as the human oncogene NET1, a Rho-GEF (Drivas, G. T. et al. (1990) Mol. Cell. Biol. -

10:1793-1798; Alberts, A. S. and Treisman, R. (1998) EMBO J. 14:4075-4085).

A novel group of GTP-binding proteins is the GTP1/OBG family, which are found in species ranging from bacteria to yeast to humans. These proteins contain characteristic GTP-binding motifs and are similar to one another but do not show sequence homology to other GTP-binding proteins.

5 The exact functions of these proteins are as yet uncertain, but they have been shown to be important for regulation of cell differentiation and development (Okamoto, S. and Ochi, K. (1998). Mol. Microbiol 30:107-119; Sazaka, T. et al. (1992) Biochem. Biophys. Res. Commun. 189:363-370).

The discovery of new GTPase associated proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis,
10 prevention, and treatment of cell proliferative, autoimmune/inflammatory, and immune system disorders.

SUMMARY OF THE INVENTION

The invention features substantially purified polypeptides, GTPase associated proteins,
15 referred to collectively as "GTPAP" and individually as "GTPAP-1," "GTPAP-2," "GTPAP-3," "GTPAP-4," "GTPAP-5," "GTPAP-6," "GTPAP-7," "GTPAP-8," "GTPAP-9," "GTPAP-10," "GTPAP-11," "GTPAP-12," "GTPAP-13," "GTPAP-14," "GTPAP-15," "GTPAP-16," "GTPAP-17," "GTPAP-18," "GTPAP-19," "GTPAP-20," "GTPAP-21," "GTPAP-22," "GTPAP-23," "GTPAP-24," "GTPAP-25," "GTPAP-26," "GTPAP-27," "GTPAP-28," and "GTPAP-29." In one aspect, the
20 invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also includes a polypeptide comprising an amino acid sequence that differs by one or more conservative amino acid substitutions from an amino acid sequence selected from the group consisting of SEQ ID NO:1-29.

The invention further provides a substantially purified variant having at least 90% amino acid
25 identity to at least one of the amino acid sequences selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also includes an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide
30 encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The

invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

The invention also provides a method for detecting a polynucleotide in a sample containing nucleic acids, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence to at least one of the polynucleotides of the sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide in the sample. In one aspect, the method further comprises amplifying the polynucleotide prior to hybridization.

The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof. The invention further provides an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof.

The invention further provides an expression vector containing at least a fragment of the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29. In another aspect, the expression vector is contained within a host cell.

The invention also provides a method for producing a polypeptide, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing a polynucleotide of the invention under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention further includes a purified antibody which binds to a polypeptide selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also provides a purified agonist and a purified antagonist to the polypeptide.

The invention also provides a method for treating or preventing a disorder associated with decreased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially

purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention also provides a method for treating or preventing a disorder associated with increased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding GTPAP.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of GTPAP.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding GTPAP were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze GTPAP, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same

meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"GTPAP" refers to the amino acid sequences of substantially purified GTPAP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of GTPAP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GTPAP either by directly interacting with GTPAP or by acting on components of the biological pathway in which GTPAP participates.

An "allelic variant" is an alternative form of the gene encoding GTPAP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding GTPAP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as GTPAP or a polypeptide with at least one functional characteristic of GTPAP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding GTPAP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding GTPAP. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent GTPAP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of GTPAP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged -

amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

5 The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein
10 molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

15 The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of GTPAP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GTPAP either by directly interacting with GTPAP or by acting on components of the biological pathway in which GTPAP participates.

20 The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind GTPAP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired.
25 Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

30 The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition containing a nucleic acid sequence which is -

complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic GTPAP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding GTPAP or fragments of GTPAP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more Incyte Clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that, when made, least interfere with the properties of the original protein, i.e., the structure and especially the function of the

protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
5	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
10	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
15	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
20	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "fragment" is a unique portion of GTPAP or the polynucleotide encoding GTPAP which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment

used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:30-58 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:30-58, for example, as distinct from any other sequence in the same genome. A fragment of SEQ ID NO:30-58 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:30-58 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:30-58 and the region of SEQ ID NO:30-58 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-29 is encoded by a fragment of SEQ ID NO:30-58. A fragment of SEQ ID NO:1-29 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-29. For example, a fragment of SEQ ID NO:1-29 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-29. The precise length of a fragment of SEQ ID NO:1-29 and the region of SEQ ID NO:1-29 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the

substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example,
5 as defined by a particular SEQ ID number, or may be measured over a shorter length, for example,
over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at
least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous
nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported
by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a
10 length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode
similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes
in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid
sequences that all encode substantially the same protein.

15 The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to
the percentage of residue matches between at least two polypeptide sequences aligned using a
standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some
alignment methods take into account conservative amino acid substitutions. Such conservative
substitutions, explained in more detail above, generally preserve the hydrophobicity and acidity at the
20 site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default
parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e
sequence alignment program (described and referenced above). For pairwise alignments of
polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap
25 penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default
residue weight table. As with polynucleotide alignments, the percent identity is reported by
CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise
comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.9
30 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

“Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term “humanized antibody” refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY;

specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C₀t or R₀t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate" refers to a change in the activity of GTPAP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of GTPAP.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Probe" refers to nucleic acid sequences encoding GTPAP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5. 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to

5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be use to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding GTPAP, or fragments thereof, or GTPAP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA,-

RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may

have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human GTPase associated proteins (GTPAP), the polynucleotides encoding GTPAP, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, autoimmune/inflammatory, and immune system disorders.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding GTPAP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each GTPAP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The Incyte clones in column 5 were used to assemble the consensus nucleotide sequence of each GTPAP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical

methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding GTPAP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:30-58 and to distinguish between SEQ ID NO:30-58 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express GTPAP as a fraction of total tissues expressing GTPAP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing GTPAP as a fraction of total tissues expressing GTPAP. Column 5 lists the vectors used to subclone each cDNA library. Of particular note is the specific expression of SEQ ID NO:43 in only one library, a human testis tissue library; the specific expression of SEQ ID NO:49 in only 4 libraries, one of which is associated with cell proliferation and 3 of which are associated with inflammation; and the specific expression of SEQ ID NO:40 in only 5 libraries, 3 of which are associated with cell proliferation and one of which is associated with inflammation.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding GTPAP were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

The invention also encompasses GTPAP variants. A preferred GTPAP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the GTPAP amino acid sequence, and which contains at least one functional or structural characteristic of GTPAP.

The invention also encompasses polynucleotides which encode GTPAP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:30-58, which encodes GTPAP.

The invention also encompasses a variant of a polynucleotide sequence encoding GTPAP. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding GTPAP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:30-58 which has at least about 70%, or alternatively at least about 90%, or even at least about

95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:30-58. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of GTPAP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding GTPAP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring GTPAP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode GTPAP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring GTPAP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding GTPAP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding GTPAP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode GTPAP and GTPAP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding GTPAP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:30-58 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment

of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Perkin-Elmer). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding GTPAP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been

size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

5 Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate
10 software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

 In another embodiment of the invention, polynucleotide sequences or fragments thereof
15 which encode GTPAP may be cloned in recombinant DNA molecules that direct expression of GTPAP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express GTPAP.

 The nucleotide sequences of the present invention can be engineered using methods generally
20 known in the art in order to alter GTPAP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction
25 sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

 In another embodiment, sequences encoding GTPAP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.)
 Alternatively, GTPAP itself or a fragment thereof may be synthesized using chemical methods. For
30 example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of GTPAP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active GTPAP, the nucleotide sequences encoding GTPAP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding GTPAP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding GTPAP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding GTPAP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding GTPAP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding GTPAP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or

tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding GTPAP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding GTPAP can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding GTPAP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of GTPAP are needed, e.g. for the production of antibodies, vectors which direct high level expression of GTPAP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of GTPAP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of GTPAP. Transcription of sequences encoding GTPAP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding GTPAP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader

sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses GTPAP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of GTPAP in cell lines is preferred. For example, sequences encoding GTPAP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *ap^r* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system.

(See, e.g., Rhodes, C.A. (1995) *Methods Mol. Biol.* 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding GTPAP is inserted within a marker gene sequence, transformed cells containing sequences encoding GTPAP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding GTPAP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding GTPAP and that express GTPAP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of GTPAP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on GTPAP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding GTPAP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding GTPAP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for

ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding GTPAP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode GTPAP may be designed to contain signal sequences which direct secretion of GTPAP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding GTPAP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric GTPAP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of GTPAP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the GTPAP encoding sequence and the heterologous protein sequence, so that GTPAP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10).

A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled GTPAP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ^{35}S -methionine.

Fragments of GTPAP may be produced not only by recombinant means, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of GTPAP may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of GTPAP and GTPase associated proteins. In addition, the expression of GTPAP is closely associated with proliferating tissues associated with cancer and fetal development, inflamed tissues, and tissues involved in the immune response. Therefore, GTPAP appears to play a role in cell proliferative, autoimmune/inflammatory, and immune system disorders. In the treatment of disorders associated with increased GTPAP expression or activity, it is desirable to decrease the expression or activity of GTPAP. In the treatment of disorders associated with decreased GTPAP expression or activity, it is desirable to increase the expression or activity of GTPAP.

Therefore, in one embodiment, GTPAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis,

autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and an immune system disorder, such as acquired immunodeficiency syndrome (AIDS), X-linked agammaglobinemia of Bruton, common variable immunodeficiency (CVI), DiGeorge's syndrome (thymic hypoplasia), thymic dysplasia, isolated IgA deficiency, severe combined immunodeficiency disease (SCID), immunodeficiency with thrombocytopenia and eczema (Wiskott-Aldrich syndrome), Chediak-Higashi syndrome, chronic granulomatous diseases, hereditary angioneurotic edema, and immunodeficiency associated with Cushing's disease, leukemias such as multiple myeloma, and lymphomas such as Hodgkin's disease.

In another embodiment, a vector capable of expressing GTPAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified GTPAP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of GTPAP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP including, but not limited to, those listed above.

In a further embodiment, an antagonist of GTPAP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GTPAP. Examples of such disorders include, but are not limited to, those cell proliferative, autoimmune/inflammatory, and immune system disorders described above. In one aspect, an antibody which specifically binds GTPAP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for

bringing a pharmaceutical agent to cells or tissues which express GTPAP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding GTPAP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GTPAP including, but not limited to, those described above.

5 In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the
10 various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of GTPAP may be produced using methods which are generally known in the art. In particular, purified GTPAP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind GTPAP. Antibodies to GTPAP may
15 also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans,
20 and others may be immunized by injection with GTPAP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in
25 humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to GTPAP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the
30 entire amino acid sequence of a small, naturally occurring molecule. Short stretches of GTPAP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to GTPAP may be prepared using any technique which provides for

the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J.*

Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and
5 Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda,
10 S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce GTPAP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

15 Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for GTPAP may also be generated.
20 For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

25 Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between GTPAP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies
30 reactive to two non-interfering GTPAP epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for GTPAP. Affinity is expressed as an

association constant, K_a , which is defined as the molar concentration of GTPAP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple GTPAP epitopes, represents the average affinity, or avidity, of the antibodies for GTPAP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular GTPAP epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the GTPAP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of GTPAP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies. Volume I: A Practical Approach, IRL Press, Washington, DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of GTPAP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding GTPAP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding GTPAP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding GTPAP. Thus, complementary molecules or fragments may be used to modulate GTPAP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding GTPAP.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides

encoding GTPAP. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

Genes encoding GTPAP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding GTPAP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding GTPAP. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding GTPAP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques

for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding GTPAP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA
5 constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase
10 linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat.
15 Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical
25 or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of GTPAP, antibodies to GTPAP, and mimetics, agonists, antagonists, or inhibitors of GTPAP. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical
30 carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial,

intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration.

Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide. Lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's

solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of GTPAP, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example GTPAP or fragments thereof, antibodies of GTPAP, and agonists, antagonists or inhibitors of GTPAP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be

determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about $0.1 \mu\text{g}$ to $100,000 \mu\text{g}$, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind GTPAP may be used for the diagnosis of disorders characterized by expression of GTPAP, or in assays to monitor patients being treated with GTPAP or agonists, antagonists, or inhibitors of GTPAP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for GTPAP include methods which utilize the antibody and a label to detect GTPAP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring GTPAP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of GTPAP expression. Normal or standard values for GTPAP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to GTPAP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of GTPAP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding GTPAP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of GTPAP may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of GTPAP, and to monitor regulation of GTPAP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding GTPAP or closely related molecules may be used to identify nucleic acid sequences which encode GTPAP. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding GTPAP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the GTPAP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:30-58 or from genomic sequences including promoters, enhancers, and introns of the GTPAP gene.

Means for producing specific hybridization probes for DNAs encoding GTPAP include the cloning of polynucleotide sequences encoding GTPAP or GTPAP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding GTPAP may be used for the diagnosis of disorders associated with expression of GTPAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and an immune system disorder, such as acquired immunodeficiency syndrome (AIDS), X-linked agammaglobulinemia of Bruton, common variable immunodeficiency (CVI), DiGeorge's syndrome (thymic hypoplasia), thymic dysplasia, isolated IgA deficiency, severe combined immunodeficiency disease (SCID), immunodeficiency with thrombocytopenia and eczema (Wiskott-Aldrich syndrome), Chediak-Higashi syndrome, chronic granulomatous diseases, hereditary angioneurotic edema, and immunodeficiency associated with Cushing's disease, leukemias such as multiple myeloma, and lymphomas such as Hodgkin's disease. The polynucleotide sequences encoding GTPAP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered GTPAP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding GTPAP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding GTPAP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding GTPAP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of GTPAP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding GTPAP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding GTPAP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a

polynucleotide encoding GTPAP, or a fragment of a polynucleotide complementary to the polynucleotide encoding GTPAP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

5 Methods which may also be used to quantify the expression of GTPAP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is
10 presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify
15 genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)
20

In another embodiment of the invention, nucleic acid sequences encoding GTPAP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence.
25 The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355; Price, C.M. (1993) *Blood Rev.* 7:127-134; and Trask, B.J. (1991) *Trends Genet.* 7:149-154.)

30 Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the

location of the gene encoding GTPAP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

5 In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides
10 valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention
15 may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

 In another embodiment of the invention, GTPAP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a
20 solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between GTPAP and the agent being tested may be measured.

 Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are
25 synthesized on a solid substrate. The test compounds are reacted with GTPAP, or fragments thereof, and washed. Bound GTPAP is then detected by methods well known in the art. Purified GTPAP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

30 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding GTPAP specifically compete with a test compound for binding GTPAP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with GTPAP.

In additional embodiments, the nucleotide sequences which encode GTPAP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. Nos. 60/109,592, 60/118,610, and 60/127,990 are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A⁺) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the

recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSORT1 plasmid (Life Technologies), or pINCY (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent E. coli cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). Electrophoretic separation of cDNA sequencing reactions and detection of labeled

polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene

families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:30-58. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding GTPAP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of GTPAP Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:30-58 were produced by extension of

an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham

Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

5 The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA
10 recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

In like manner, the nucleotide sequences of SEQ ID NO:30-58 are used to obtain 5'
15 regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:30-58 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base
20 pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 µCi of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a
25 SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon
30 membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and

compared.

VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, supra.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the GTPAP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring GTPAP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of GTPAP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the GTPAP-encoding transcript.

IX. Expression of GTPAP

Expression and purification of GTPAP is achieved using bacterial or virus-based expression

systems. For expression of GTPAP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express GTPAP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of GTPAP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding GTPAP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, GTPAP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from GTPAP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified GTPAP obtained by these methods can be used directly in the following activity assay.

X. Demonstration of GTPAP Activity

The role of GTPAP can be assayed in vitro by monitoring the mobilization of Ca^{++} as part of the signal transduction pathway. (See, e.g., Grynkievicz, G. et al. (1985) J. Biol. Chem. 260:3440; McColl, S. et al. (1993) J. Immunol. 150:4550-4555; and Aussel, C. et al. (1988) J. Immunol. 140:215.) The assay requires preloading neutrophils or T cells with a fluorescent dye such as FURA-2.

Upon binding Ca^{++} , FURA-2 exhibits an absorption shift that can be observed by scanning the excitation spectrum between 300 and 400 nm, while monitoring the emission at 510 nm. When the cells are exposed to one or more activating stimuli artificially (i.e., anti-CD3 antibody ligation of the T cell receptor) or physiologically (i.e., by allogeneic stimulation), Ca^{++} flux takes place. Ca^{++} flux results from the release of Ca^{++} from intracellular organelles or from Ca^{++} entry into the cell through activated Ca^{++} channels. This flux can be observed and quantified by assaying the cells in a fluorometer or fluorescence activated cell sorter. Measurements of Ca^{++} flux are compared between cells in their normal state and those preloaded with GTPAP. Increased mobilization attributable to increased GTPAP availability results in increased emission.

Alternatively, GTPAP activity is measured by quantifying the amount of a non-hydrolyzable GTP analogue, GTP γ S, bound over a 10 minute incubation period. Varying amounts of GTPAP are incubated at 30°C in 50mM Tris buffer, pH 7.5, containing 1mM dithiothreitol, 1mM EDTA and 1 μ M [35 S]GTP γ S. Samples are passed through nitrocellulose filters and washed twice with a buffer consisting of 50mM Tris-HCl, pH 7.8, 1mM NaN_3 , 10mM MgCl_2 , 1mM EDTA, 0.5mM dithiothreitol, 0.01mM PMSF, and 200mM NaCl. The filter-bound counts are measured by liquid scintillation to quantify the amount of bound [35 S]GTP γ S. GTPAP activity may also be measured as the amount of GTP hydrolysed over a 10 minute incubation period at 37°C. GTPAP is incubated in 50mM Tris-HCl buffer, pH 7.8, containing 1mM dithiothreitol, 2mM EDTA, 10 μ M [α - 32 P]GTP, and 1 μ M H-rab protein. GTPase activity is initiated by adding MgCl_2 to a final concentration of 10 mM. Samples are removed at various time points, mixed with an equal volume of ice-cold 0.5mM EDTA, and frozen. Aliquots are spotted onto polyethyleneimine-cellulose thin layer chromatography plates, which are developed in 1M LiCl, dried, and autoradiographed. The signal detected is proportional to GTPAP activity.

Alternatively, GTPAP activity may be demonstrated as the ability to interact with its associated $\text{G}\alpha$ or LMW GTPase in an in vitro binding assay. The candidate GTPases are expressed as fusion proteins with glutathione S-transferase (GST), and purified by affinity chromatography on glutathione-Sepharose. The GTPases are loaded with GDP by incubating 20 mM Tris buffer, pH 8.0, containing 100 mM NaCl, 2 mM EDTA, 5 mM MgCl_2 , 0.2 mM DTT, 100 μ M AMP-PNP and 10 μ M GDP at 30°C for 20 minutes. GTPAP is expressed as a FLAG fusion proteins in a baculovirus system. Extracts of these baculovirus cells containing GTPAP-FLAG fusion proteins are precleared with GST beads, then incubated with GST-GTPase fusion proteins. The complexes formed are precipitated by glutathione-Sepharose and separated by SDS-polyacrylamide gel electrophoresis. The separated proteins are blotted onto nitrocellulose membranes and probed with commercially available anti-

FLAG antibodies. GTPAP activity is proportional to the amount of GTPAP-FLAG fusion protein detected in the complex.

XI. Functional Assays

GTPAP function is assessed by expressing the sequences encoding GTPAP at

5 physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT (Life Technologies) and pCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell
10 line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM),
15 an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and
20 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

25 The influence of GTPAP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding GTPAP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake
30 Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding GTPAP and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of GTPAP Specific Antibodies

GTPAP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the GTPAP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-GTPAP activity by, for example, binding the peptide or GTPAP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIII. Purification of Naturally Occurring GTPAP Using Specific Antibodies

Naturally occurring or recombinant GTPAP is substantially purified by immunoaffinity chromatography using antibodies specific for GTPAP. An immunoaffinity column is constructed by covalently coupling anti-GTPAP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing GTPAP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of GTPAP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/GTPAP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and GTPAP is collected.

XIV. Identification of Molecules Which Interact with GTPAP

GTPAP, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled GTPAP, washed, and any wells with labeled GTPAP complex are assayed. Data obtained using different concentrations of GTPAP are used to calculate values for the number, affinity, and association of

GTPAP with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

1.02257 15.2201 0986679 15.2201 0986679

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	30	708398	SYNORAT04	568987X31 (MMLR3DT01), 708398H1, 708398X11, 708398X15, 708398X16, 708398X17, and 708398X21 (SYNORAT04), 2170523F6 (ENDCNOT03), 3374750H1 (CONNTUT05)
2	31	1259937	MENITUT03	913652R6 (STOMNOT02), 1259937F6 and 1259937H1 (MENITUT03), 1476721F1 (CORPNOT02), 1729248F6 (BRSTTUT08), 2191963H1 (THYRTUT03), 3129757F6 (LUNGUTUT12), 3268746X15F1 (BRAINTUT02), 3428891F6 (SKINNOT04)
3	32	1452285	PENITUT01	1452285F6 and 1452285H1 (PENITUT01), 2605011H1 (LUNGUTUT07), 3505135H1 (ADRENOT11)
4	33	1812894	PROSTUT12	1812894H1, 1812894X12 and 1809113T6 (PROSTUT12), 1904479F6 (OVARNOT07), 2232535X15F1 and 2232535X18F1 (PROSNOT16), 2267486X16C1 (UTRSNOT02), 2508562F6 (CONUTUT01)
5	34	3074884	BONEUNT01	225362F1 (PANCNOT01), 900707R1 (BRSTTUT03), 1339234F6 (COLNTUT03), 1759046R6 (PITUNOT03), 3074884H1 (BONEUNT01), SBDA02767F1
6	35	3452277	UTRSNON03	1684553F6 (PROSNOT15), 1951534H1 (PITUNOT01), 3452277H1 (UTRSNON03), 4092781T6 (BSCNSZT01), SBFA01413F1, SBFA03044F1, SBFA01805F1
7	36	4203832	BRAITUT29	723394F1 (SYNOOAT01), 8622290R1, and 8622290T1 (BRAITUT03), 1560918F1 (SPLNNOT04), 3509241H1 (CONCNOT01), 4203832H1 (BRAITUT29)
8	37	104368	BMARNOT02	104368H1 (BMARNOT02), SAEA03574F1, SAEA01063F1, SAEA00392F1, SAEA02287F1
9	38	1441680	THYRNOT03	1441680F6, 1441680H1, and 1441680T6 (THYRNOT03), 1904222F6 (OVARNOT07), 2477983F6 (SMCANOT01)
10	39	1494955	PROSNON01	965986R1 (BRSTNOT05), 1429037F1 and 1429037T1 (SINTBST01), 1453487F6 (PENITUT01), 1486114H1 (CORPNOT02), 1494955H1 (PROSNON01), 1995426R6 (BRSTTUT03), 2112074X18F1 and 2112348R6 (BRAITUT03)
11	40	1508161	LUNGNOT14	1508161F6 and 1508161H1 (LUNGNOT14), 3334303H1 (BRAIFET01), 4755656H1 (BRAHNOT01)
12	41	1811877	PROSTUT12	493795H1 (HNT2NOT01), 1573136H1 (LNODNOT03), 1811877F6 and 1811877H1 (PROSTUT12), 1825223F6 (LSUBNOT03), 2454143H1 (ENDANOT01), 2651022H1 (BLADTUT08), 3487062H1 (EPIGNOT01), 4536531H1 (OVARNOT12), 4795253H1 (LIVRTUT09), 4854087H1 (TESTNOT10), 4906149H2 (TLYMNOT08), 5196386H1 (LUNLTUT04)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
13	42	1848674	LUNGFET03	1574127F6, 3857867X306F1, and 3857867X313F1 (LNODNOT03), 1848674H1 (LUNGFET03), 1877170F6 (LEUKNOT03), 2695307H1 (UTRSNOT12), 4148654H1 (SINITUT04), 4984182H1 (HELATXT05), 5288671H1 (LIVRTUS02)
14	43	2012970	TESTNOT03	2012970H1, 2012970R6, 2012970X11F (TESTNOT03)
15	44	2254315	OVARTUT01	022341F1 (ADENINB01), 198476R6 (KIDNNOT02), 2254315H1 (OVARTUT01), 2370170F6 (ADRENOT07), 2451278F6 (ENDANOT01)
16	45	2415545	HNT3AZT01	775722H1 (COLNNOT05), 870320R6 (LUNGAST01), 889023R1 (STOMTUT01), 895724R1 (BRSTNOT05), 1398541F1 (BRAITUT08), 1662585F6 (BRSTNOT09), 2415545H1 (HNT3AZT01), 2985066H1 (CARGDIT01), 3462702H1 (293TF2T01)
17	46	2707969	PONSATZ01	282552R1, 282552X23, and 282552X7 (CARDNOT01), 889783R1 (STOMTUT01), 1995451R6 (BRSTTUT03), 2707969H1 (PONSATZ01), SAAC00359R1.comp, SAAB00136R1, SAAC00330R1
18	47	2817769	BRSTNOT14	041660R1 (TBLYNOT01), 077378R1 (SYNORAB01), 740028R1 (PANCNOT04), 1593201F6 (BRAINOT14), 1924025R6 (BRSTTUT01), 2817769H1 (BRSTNOT14)
19	48	2917557	THYMFET03	473002F1 and 473002R1 (MMLR1DT01), 690999R6 (LUNGUT02), 997483R1 (KIDNTUT01), 1430662F6 (SINTBST01), 1514017F1 (PANCNTUT01), 1740475R6 (HIPONON01), 2109547H1 (BRAITUT03), 2917557H1 (THYMFET03), 4309528H1 (BRAUNOT01), 4990135H1 (LIVRTUT11)
20	49	3421335	UCMCNOT04	777588R6 and 777588T6 (COLNNOT05), 3421335H1 (UCMCNOT04)
21	50	605761	BRSTTUT01	605761F1, 605761H1, and 605761R6 (BRSTTUT01), 1271131X15 (TESTTUT02), 1516985F1 (PANCNTUT01), 1524935H1 (UCMCL5T01), 2234846F6 (PANCNTUT02)
22	51	483862	HNT2RAT01	483862H1 and 483862R1 (HNT2RAT01), 1750781X305F1, 1750781X307D2 (LIVRTUT01)
23	52	1256777	MENITUT03	264041R6 (HNT2AGT01), 826449R1 (PROSNOT06), 1256777H1 (MENITUT03), 2276061R6 (PROSNON01), 4614049H1 (BRAHNOT01)
24	53	2198779	SPLNFET02	1557708F6 (BLADTUT04), 1922490R6 (BRSTTUT01), 2198779H1 (SPLNFET02), 2541193F7 (BONRTUT01), 3039254F6 (BRSTNOT16), 3057079H1 (LNODNOT08), 3105017H1 (COLNUCT03), 4239592H1 (SYNWDIT01), 5064513H1 (ARTFTDT01)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
25	54	2226116	SEMVNOT01	1662607F6 (BRSTNOT09), 1662607T6 (BRSTNOT09), 2226116F6 (SEMVNOT01), 2226116H1 (SEMVNOT01), 2930011F6 (TLYMNOT04), 3015747T6 (MUSCNOT07), 4087670H1 (LIVRNOT06)
26	55	2504472	CONUTUT01	420365F1 (BRSTNOT01), 762246R1 (BRAITUT02), 907754R2 (COLNNOT09), 1007508H1 (HEALDIT02), 1302342F6 (PLACNOT02), 1913887H1 (PROSTUT04), 2023822F6 (CONNNOT01), 2023822X11R1 (CONNNOT01), 2504472H1 (CONUTUT01), 2951618F6 (KIDNFET01)
27	56	3029920	HEARFET02	354846T6 (RATRNUT01), 418533R6 (BRSTNOT01), 935073R1 (CERVNOT01), 1340722F1 (COLNTUT03), 1416203T6 (BRAINOT12), 1524567F1 (UCMCL5T01), 1773043H1 (MENTUNON3), 2590310H2 (LUNGNOT22), 3029920H1 (HEARFET02), 4873053H1 (COLDNOT01), 5687696H1 (BRAIUNT01)
28	57	3332415	BRAIFET01	118166R1 (MUSCNOT01), 1257348H1 (MENITUT03), 1288237T6 (BRAINOT11), 1335936F6 (COLNNOT13), 1452268H1 (PENITUT01), 1996016R6 (BRSTTUT03), 2116665R6 (BRSTTUT02), 2206894F6 (SINTFET03), 2540063H1 (BONRTUT01), 2808268H1 (BLADTUT08), 3086221H1 (HEAONOT03), 3127508H1 (LUNGTUT12), 3295812H1 (TLYJINT01), 3332415H1 (BRAIFET01), 3604705H1 (LUNGNOT30), 4821203H1 (PROSTUT17), 4970353H1 (KIDEUNC10), 5055775H1 (COLATMT01)
29	58	4031536	BRAINOT23	029167X3 (SPLNFET01), 350137R1 (LVENNOT01), 408825X1 (EOSIHET02), 689446X23 (LUNGTUT02), 1963062R6 (BRSTNOT04), 2288043R6 (BRAINON01), 4031536H1 (BRAINOT23)

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
1	1002	T30 S224 T405 S499 T533 S558 S701 T737 T845 S864 S6 T152 T268 T412 T442 T464 T514 T528 T693 S814 S815 S823 T880 Y117 Y842	N446	G524-T531: ATP/GTP- binding site motif	GTP-binding protein [Mus musculus] g53169	BLAST MOTIFS
2	338	S21 S77 T86 S200 T246 T299 S77 S306 Y131	N244		cAMP- regulated Guanine nucleotide exchange factor [Rattus norvegicus] g4079657	BLAST
3	211	S159 S199	N33 N74	G16-T23: ATP/GTP- binding site motif	GTP-binding protein [Rattus norvegicus] g206543	BLAST MOTIFS PFAM BLOCKS PRINTS
4	516	T14 S42 T237 S270 S347 S360 T371 T395 T433 S500 T3 S13 S96 T316 S430			Fos-related antigen [Rattus norvegicus] g1016712 Rabaptin-4 [H. sapiens] g3832516	BLAST MOTIFS
5	445	T44 T114 T219 T297 S314 S341 S356 T412 T24 S72 T91 T328 T388 T394		G230-T237: ATP/GTP- binding site motif	GTP-binding protein [H. sapiens] g2765411	BLAST MOTIFS

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
6	445	S174 S202 S289 S29 S305 S323 T434 T11 T147 T197 T198 S270 S273 S371 S397 Y125	N73		Regulator of G-protein signaling-9 [H. sapiens] g3284012	BLAST
7	281	S182 S210 S254 S13 T56 S110 S182 S32 T46 S66 S177	N130 N181	G31-T38:ATP/GTP- binding site motif	Putative ras- like protein [H. sapiens] g4092830	MOTIFS PRINTS BLAST PFAM
8	301	S92 T2 T3 Y15 S18 S19 S20 S25 S97 T120 S165 S296 T94 S116 T120 S284		E47-G66, S116-E178, Y188-G272: Phosducin signature	Phosducin- like protein [Rattus rattus] g1323727	MOTIFS BLAST PRINTS
9	485	T6 Y57 S82 T91 S112 S187 T231 T257 S309 T6 T81 S132 S157 S210 S241 T462	N460	L49-S82: Beta G protein	Similar to WD domain Beta transducin- like protein [C. elegans] g5596646	MOTIFS BLAST PRINTS
10	447	S420 S94 T107 S118 T167 T179 T308 S390 S39 S58 T78 T113 S129 T160 T167 Y174 T199 S216 S291 T302 T323 T359 T384 S423 T438	N76 N92 N231 N289 N378 N421	M294-T308: Beta transducin	WS beta- transducin repeat protein [Homo sapiens] g4704417	MOTIFS BLAST
11	199	S90 T55 T140 S190		K6-E130: Ras Guanine exchange factor	Putative guanine nucleotide releasing factor [Drosophila affinis] g2981229	MOTIFS BLAST PFAM

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
12	694	S57 S67 S99 T150 T346 S416 S467 S500 T522 T684 S99 T156 S209 S285 T331 T360 T388 T430 T477 T650 T688		L10-I24, M96-L110: Beta transducin	Transducin- like protein [H. sapiens] g414536	MOTIFS BLAST
13	654	T10 S15 T49 S97 S102 S104 S112 S113 S377 S432 S638 T46 S54 S84 S97 T177 S217 T307 S401 S450 S504 T515 S546 T547 S561 Y618	N353 N362 N502	L197-F211: Beta transducin	Similar to the beta transducin family [C. elegans] g2315521	MOTIFS BLAST
14	180	S14		G23-S30: ATP-GTP binding site	Rab7C (small GTP binding protein) [Lotus japonicus] g1370186	MOTIFS BLAST
15	374	T100 T249 S260 T308 T328 S338 S351 S30 T73 T157 S237 T308	N114 N189 N222	G26-T33: ATP-GTP binding site	ATP (GTP)- binding protein [H. sapiens] g3646130	MOTIFS BLAST
16	649	S67 T344 S366 S63 S68 S75 S122 S177 S265 T282 T332 S373 S380 S563 T569 S634 S20 T94 S128 S314 T382 T385 T458 T559		F307-S544: Probable rabGAP domain	Similar to probable rabGAP [C. elegans] g3925265	MOTIFS BLAST PFAM
17	698	T244 S262 S17 T41 T42 T196 S206 S317 S479 S522 S556 T586 T680 T31 S95 T99 T140 T173 S257 T322 S374 T450 S568 T619	N171 N194 N685		Small Gtp- binding protein associated protein [Mus musculus] g725274	MOTIFS BLAST

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
18	396	T325 S115 T133 S232 S275 T336 S22 T221 S232 T320	N60 N230 N286	G29-S36: ATP-GTP binding site	Putative GTP- binding protein [C. elegans] g3880615	MOTIFS BLAST
19	634	T197 S3 S5 S9 T14 S132 T197 T285 T553 T40 T56 S160 T189 S261 S582 Y20 Y396 Y419		G52-T59: ATP-GTP binding site	Putative GTP- binding protein [H. sapiens] g3169010	MOTIFS BLAST
20	196	T60 S73 S90 S99 S73 S193		G19-T26: ATP-GTP binding site	Kidney injury associated protein HW052 Acc No W86322 ADP- ribosylation factor-like protein 3 [Rattus norvegicus] g560006	MOTIFS BLAST
21	446	T10 T24 T93 S122 T243 S263 S270 T305 S317 S325 T357 S372 T379 S100 S170 S223 T227 S285 T348	N79	L323-L337: Beta transducin	Putative WD40 repeat protein [A. thaliana] g4191784	MOTIFS BLAST
22	265	T184 T76 T137 S139 T161 T174 T183 S213	N159	L141, L148, L155 L: zipper gene regulatory motif	TipD; similar to beta transducin family [D. discoideum] g2407788	MOTIFS BLAST

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
23	185	T55 S111 S127 S148 S171 S14 S94 Y103		G10-T17: ATP/GTP binding site (P- loop) A4-S72: Ras domain		MOTIFS PFAM PRINTS
24	554	S388 T488 S30 S75 T111 S149 S220 S237 T255 S305 S325 T339 T359 S363 S509 S172 T195 S211 T378 T438 T470 Y203	N5	N297-D336, P345- D383, G481-Q519: Beta-transducin WD40 repeats	WD-repeat protein [Arabidopsis thaliana] g3924603	BLAST MOTIFS PFAM PRINTS
25	434	S164 S341 T347 S36 S68 S92 T286 S364	N22 N383	G259-S266:ATP/GTP binding site (P- loop): G113-R433: GTP1/OBG domain	Predicted GTP binding protein [C. elegans] g3878629	BLAST MOTIFS PFAM BLOCKS PRINTS
26	826	S122 T243 T247 T427 S454 S519 T528 S623 S701 S715 S809 T58 S143 S266 T411 S505 S577 S603 T661 S735 T753 S791 T815	N23 N264 N576 N600 N789	R48-E91, L97-S143, F197 K237, V273- W319, W378-A416, W604 K642, A659- G697: Beta- transducin WD40 repeats	Predicted WD repeat protein [S. cerevesiae] P42935	BLAST MOTIFS PFAM PRINTS
27	618	T414 S59 T105 S126 T139 T143 S196 T203 S311 S325 T370 T390 S477 T483 S541 T583 T94 S148 T247 Y160 Y383 Y456	N118 N154 N346	G11-T18, G425-S432: ATP/GTP binding site (P-loop) R6-K187: Ras domain	GTP-binding protein APD08 [H.sapiens] Accession W75771	BLAST MOTIFS PFAM PRINTS

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
28	596	S17 S21 S50 S152 S153 T533 S539 T594 S36 S38 S80 T163 T169 S183 S211 T240 S306 T329 T417 S457 S508 T545 S45 T64 S88 T124 S139 S299 S451 S459 S528 S568 Y180 Y364		A178-L355: Rho- family guanine nucleotide exchange factor (RhoGEF) domain	Guanine nucleotide regulatory protein (NET1 homologue) [Mus musculus] g3834631	BLAST MOTIFS PFAM BLOCKS
29	589	T108 S20 T90 S127 S176 S278 S467 T521 S522 T189 S254 T284 T292 T321 T324 T345 T364 T423 S444 T484 T527	N572	L252-S289, G293- N329, G333-D369, G373-D409, E413- D449, G453-D489, G493-D532: Beta- transducin WD40 repeats R160-K206: F-box domain	SEL-10 [C.elegans] g2677836	BLAST MOTIFS PFAM PRINTS

Table 3

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
30	628-711	Reproductive (0.256) Nervous (0.154) Gastrointestinal (0.154)	Cell Proliferation (0.692) Inflammation (0.372)	PSPORT1
31	1094-1129	Reproductive (0.268) Cardiovascular (0.146) Nervous (0.146)	Cell Proliferation (0.731) Inflammation (0.219) Neurological (0.049)	pINCY
32	652-703	Cardiovascular (0.375) Reproductive (0.375) Dermatologic (0.125) Endocrine (0.125)	Cell Proliferation (0.875) Trauma (0.125)	pINCY
33	1224-1292	Reproductive (0.412) Gastrointestinal (0.147) Hematopoietic/Immune (0.147)	Cell Proliferation (0.647) Inflammation (0.264)	pINCY
34	16-65	Nervous (0.211) Reproductive (0.197) Gastrointestinal (0.169)	Cell Proliferation (0.507) Inflammation (0.352)	pINCY
35	947-1043	Reproductive (0.444) Nervous (0.333) Gastrointestinal (0.111) Urologic (0.111)	Cell Proliferation (0.667) Inflammation (0.111) Neurological (0.111)	pINCY
36	840-1001	Nervous (0.340) Reproductive (0.208) Gastrointestinal (0.151)	Cell Proliferation (0.641) Inflammation (0.302) Neurological (0.038)	pINCY

Table 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
37	507-551	Hematopoietic/Immune (0.269) Nervous (0.269) Reproductive (0.154)	Inflammation (0.423) Cell Proliferation (0.269)	PBLUESCRIPT
38	218-262	Cardiovascular (0.357) Nervous (0.214) Gastrointestinal (0.143)	Cell Proliferation (0.572) Inflammation (0.214)	pINCY
39	164-208	Nervous (0.280) Reproductive (0.260) Developmental (0.120)	Cell Proliferation (0.740) Inflammation (0.180)	PSPORT1
40	369-411	Cardiovascular (0.250) Developmental (0.250) Gastrointestinal (0.250)	Cell Proliferation (0.500) Inflammation (0.250)	pINCY
41	272-316	Reproductive (0.392) Gastrointestinal (0.118) Hematopoietic/Immune (0.118)	Cell Proliferation (0.626) Inflammation (0.137)	pINCY
42	664-708	Nervous (0.211) Reproductive (0.211) Cardiovascular (0.158)	Cell Proliferation (0.614) Inflammation (0.281)	pINCY
43	226-270	Reproductive (1.000)	Inflammation (1.000)	PBLUESCRIPT
44	11-55	Reproductive (0.254) Gastrointestinal (0.206) Cardiovascular (0.159)	Cell Proliferation (0.698) Inflammation (0.206)	PSPORT1

Table 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
45	637-681	Reproductive (0.281) Nervous (0.188) Gastrointestinal (0.156)	Cell Proliferation (0.781) Inflammation (0.234)	pINCY
46	1016-1060	Nervous (0.330) Reproductive (0.183) Hematopoietic/Immune (0.122)	Cell Proliferation (0.582) Inflammation (0.235)	pINCY
47	737-781	Nervous (0.218) Reproductive (0.188) Gastrointestinal (0.158)	Cell Proliferation (0.655) Inflammation (0.211)	pINCY
48	469-513	Reproductive (0.222) Hematopoietic/Immune (0.160) Nervous (0.160)	Cell Proliferation (0.543) Inflammation (0.272)	pINCY
49	226-270	Gastrointestinal (0.333) Hematopoietic/Immune (0.333) Reproductive (0.333)	Inflammation (1.000)	pINCY
50	456-500	Reproductive (0.289) Gastrointestinal (0.133) Hematopoietic/Immune (0.133)	Cell Proliferation (0.778) Inflammation (0.156)	PSPORT1
51	252-296	Nervous (0.500) Gastrointestinal (0.200) Cardiovascular (0.100)	Cell Proliferation (1.000) Inflammation (0.200)	PBLUESCRIPT
52	60-104	Nervous (0.326) Reproductive (0.326) Cardiovascular (0.152)	Cell proliferation (0.565) Inflammation (0.369)	pINCY

Table 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
53	488-532	Reproductive (0.232) Nervous (0.195) Hematopoietic/Immune (0.146)	Cell proliferation (0.622) Inflammation (0.427)	pINCY
54	686-730	Reproductive (0.250) Gastrointestinal (0.150) Hematopoietic/Immune (0.150)	Cell proliferation (0.700) Inflammation (0.400)	pINCY
55	543-587 1299-1343	Reproductive (0.282) Nervous (0.155) Gastrointestinal (0.146)	Cell proliferation (0.592) Inflammation (0.359)	pINCY
56	345-389 792-836	Nervous (0.268) Reproductive (0.169) Cardiovascular (0.113) Hematopoietic/Immune (0.113)	Cell proliferation (0.606) Inflammation (0.296)	pINCY
57	163-207	Reproductive (0.270) Gastrointestinal (0.189) Nervous (0.156)	Cell proliferation (0.705) Inflammation (0.254)	pINCY
58	381-425 726-770	Nervous (0.317) Reproductive (0.250) Gastrointestinal (0.117)	Cell proliferation (0.450) Inflammation (0.283)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
30	SYNORAT04	This library was constructed using RNA isolated from the wrist synovial membrane tissue of a 62-year-old female with rheumatoid arthritis.
31	MENITUT03	This library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old female during excision of a cerebral meningeal lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
32	PENITUT01	This library was constructed using RNA isolated from tumor tissue removed from the penis of a 64-year-old male during penile amputation. Pathology indicated a fungating invasive grade 4 squamous cell carcinoma involving the inner wall of the foreskin and extending onto the glans penis. Patient history included benign neoplasm of the large bowel, atherosclerotic coronary artery disease, angina pectoris, gout, and obesity. Family history included malignant pharyngeal neoplasm, chronic lymphocytic leukemia, and chronic liver disease.
33	PROSTUT12	This library was constructed using RNA isolated from prostate tumor tissue removed from a 65-year-old male during a radical prostatectomy. Pathology indicated an adenocarcinoma (Gleason grade 2+2). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA).
34	BONEUNT01	This library was constructed using RNA isolated from Saos-2, a primary osteogenic sarcoma cell line (ATCC HTB-85) derived from an 11-year-old Caucasian female.
35	UTRSNON03	This library was constructed from 6.4 million independent clones from a uterine library. RNA for these libraries was isolated from uterine myometrial tissue removed from a 41-year-old female during a vaginal hysterectomy with dilation and curettage. The endometrium was secretory and contained fragments of endometrial polyps. Benign endo- and ectocervical mucosa were identified in the endocervix. Pathology for the associated tumor tissue indicated uterine leiomyoma. The normalization and hybridization conditions were adapted from Soares et al. (Proc.Natl.Acad.Sci. USA (1994) 91:9928).
36	BRAITUT29	This library was constructed using RNA isolated from brain tumor tissue removed from the parietal lobe of a 43-year-old female during excision of a cerebral meningeal lesion. Pathology indicated high grade glioma. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, benign hypertension, and hyperlipidemia.
37	BMARNOT02	This library was constructed using RNA isolated from the bone marrow of 24 male and female Caucasian donors, 16 to 70 years old. (RNA came from Clontech.)

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
38	THYRN0T03	This library was constructed using RNA isolated from thyroid tissue removed from the left thyroid of a 28-year-old Caucasian female during a complete thyroidectomy. Pathology indicated a small nodule of adenomatous hyperplasia present in the left thyroid. Pathology for the associated tumor tissue indicated dominant follicular adenoma, forming a well-encapsulated mass in the left thyroid.
39	PROSN0N01	This normalized library was constructed from 4.4 million independent clones from a prostate library. Starting RNA was made from prostate tissue removed from a 28-year-old Caucasian male who died from a self-inflicted gunshot wound. The normalization and hybridization conditions were adapted from Soares, M.B. et al. (1994) Proc. Natl. Acad. Sci. USA 91:9228-9232, using a longer (19 hour) reannealing hybridization period.
40	LUNGNOT14	This library was constructed using RNA isolated from lung tissue removed from the left lower lobe of a 47-year-old Caucasian male during a segmental lung resection. Pathology for the associated tumor tissue indicated a grade 4 adenocarcinoma, and the parenchyma showed calcified granuloma. Patient history included benign hypertension and chronic obstructive pulmonary disease. Family history included type II diabetes and acute myocardial infarction.
41	PROSTUT12	This library was constructed using RNA isolated from prostate tumor tissue removed from a 65-year-old Caucasian male during a radical prostatectomy. Pathology indicated an adenocarcinoma (Gleason grade 2+2). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA).
42	LUNGFET03	This library was constructed using RNA isolated from lung tissue removed from a Caucasian female fetus who died at 20 weeks' gestation.
43	TESTNOT03	This library was constructed using RNA isolated from testicular tissue removed from a 37-year-old Caucasian male, who died from liver disease. Patient history included cirrhosis, jaundice, and liver failure.
44	OVARTUT01	This library was constructed using RNA isolated from ovarian tumor tissue removed from a 43-year-old Caucasian female during removal of the fallopian tubes and ovaries. Pathology indicated grade 2 mucinous cystadenocarcinoma involving the entire left ovary. Patient history included mitral valve disorder, pneumonia, and viral hepatitis. Family history included atherosclerotic coronary artery disease, pancreatic cancer, stress reaction, cerebrovascular disease, breast cancer, and uterine cancer.
45	HNT3AZT01	This library was constructed using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated for three days with 0.35 micromolar 5-aza-2'-deoxycytidine (AZ).

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
46	PONSAZT01	This library was constructed using RNA isolated from diseased pons tissue from the brain of a 74-year-old Caucasian male who died from Alzheimer's disease.
47	BRSTNOT14	This library was constructed using RNA isolated from breast tissue obtained from a 62-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated an invasive grade 3 (of 4), nuclear grade 3 (of 3) adenocarcinoma, ductal type. Patient history included a benign colon neoplasm, hyperlipidemia, cardiac dysrhythmia, and obesity. Family history included atherosclerotic coronary artery disease, myocardial infarction, colon cancer, ovarian cancer, lung cancer, and cerebrovascular disease.
48	THYMFET03	This library was constructed using RNA isolated from thymus tissue removed from a Caucasian male fetus.
49	UCMCNOT04	This library was constructed using RNA isolated from mononuclear cells obtained from the umbilical cord blood of multiple individuals of mixed age and sex. The cells were treated with G-CSF.
50	BRSTTUT01	This library was constructed using RNA isolated from breast tumor tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated invasive grade 4 mammary adenocarcinoma of mixed lobular and ductal type, extensively involving the left breast. Family history included benign hypertension, atherosclerotic coronary artery disease, cerebrovascular disease, and depressive disorder.
51	HNT2RAT01	This library was constructed at Stratagene (STR937231), using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated with retinoic acid for 24 hours.
52	MENITUT03	This library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
53	SPLNFET02	This library was constructed using RNA isolated from spleen tissue removed from a Caucasian male fetus, who died at 23 weeks' gestation.
54	SEMVNOT01	This library was constructed using RNA isolated from seminal vesicle tissue removed from a 58-year-old Caucasian male during radical prostatectomy. Pathology for the associated tumor tissue indicated adenocarcinoma (Gleason grade 3+2) of the prostate. Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA). Family history included a malignant breast neoplasm.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
55	CONUTUT01	This library was constructed using RNA isolated from sigmoid mesentery tumor tissue obtained from a 61-year-old female during a total abdominal hysterectomy and bilateral salpingo-oophorectomy with regional lymph node excision. Pathology indicated a metastatic grade 4 malignant mixed mullerian tumor present in the sigmoid mesentery at two sites.
56	HEARFET02	This library was constructed using RNA isolated from heart tissue removed from a Caucasian male fetus, who was stillborn at 23 weeks' gestation with a hypoplastic left heart.
57	BRAIFET01	This library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus, who was stillborn at 23 weeks' gestation with a hypoplastic left heart.
58	BRAINOT23	This library was constructed using RNA isolated from right temporal lobe tissue removed from a 45-year-old Black male during a brain lobectomy. Pathology for the associated tumor tissue indicated dysembryoplastic neuroepithelial tumor of the right temporal lobe. The right temporal region dura was consistent with calcifying pseudotumor of the neuraxis. The patient presented with convulsive intractable epilepsy, partial epilepsy, and memory disturbance. Patient history included obesity, meningitis, backache, unspecified sleep apnea, acute stress reaction, acquired knee deformity, and chronic sinusitis. Family history included obesity, benign hypertension, cirrhosis of the liver, alcohol abuse, hyperlipidemia, cerebrovascular disease, and type II diabetes.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value= 1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLOCKS IMPROVED Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality score > GCG-specified "HIGH" value for that particular Prosite motif. Generally, score = 1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score = 120 or greater; Match length = 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score = 3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <i>supra</i> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	